

SOME UNIFIED INTEGRALS ASSOCIATED WITH GENERALIZED BESSEL-MAITLAND FUNCTION

M.S. ABOUZAIID¹, A.H. ABUSUFIAN², K.S. NISAR^{2,*}

ABSTRACT. Generalized integral formulas involving the generalized Bessel-Maitland function are considered and it expressed in terms of generalized Wright hypergeometric functions. By assuming appropriate values of the parameters in the main results, we obtained some interesting results of ordinary Bessel function.

1. INTRODUCTION AND PRELIMINARIES

The Bessel-Maitland $J_v^\alpha(z)$ defined by the following series representation [1]

$$J_v^\delta(z) = \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma(\delta n + v + 1)} \quad (1.1)$$

The applications of Bessel-Maitland function are found in the field of applied science, engineering, biological, chemical and physical science is found in the book of Watson [15]. The generalized Bessel-Maitland function investigated and studied in [25] and defined it as:

$$J_{v,q}^{\delta,\gamma}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn} (-z)^n}{n! \Gamma(\delta n + v + 1)} \quad (1.2)$$

where $\mu, v, \gamma \in C$; $\Re(\mu) \geq 0$, $\Re(v) > -1$, $\Re(\gamma) \geq 0$, $q \in (0, 1) \cup \mathbb{N}$ and

$$(\gamma)_0 = 1, (\gamma)_{q,n} = \frac{\Gamma(\gamma + qn)}{\Gamma(\gamma)} \quad (1.3)$$

The generalized hypergeometric function represented as follows [24]:

$${}_pF_q \left[\begin{matrix} (\alpha_p); \\ (\beta_q); \end{matrix} z \right] = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^p (\alpha_j)_n}{\prod_{j=1}^q (\beta_j)_n} \frac{z^n}{n!}, \quad (1.4)$$

provided $p \leq q$; $p = q + 1$ and $|z| < 1$

2010 *Mathematics Subject Classification.* 33C10, 33C20, 26A33, 33B15.

Key words and phrases. Bessel-Maitland function; Generalized Wright hypergeometric function; Gamma function; Integral formula.

* Corresponding author.

where $(\lambda)_n$ is well known Pochhammer symbol defined for (for $\lambda \in \mathbb{C}$) (see [24])

$$(\lambda)_n := \begin{cases} 1 & (n = 0) \\ \lambda(\lambda + 1) \dots (\lambda + n - 1) & (n \in \mathbb{N} := \{1, 2, 3, \dots\}) \end{cases} \quad (1.5)$$

$$(\lambda)_n = \frac{\Gamma(\lambda + n)}{\Gamma(\lambda)} \quad (\lambda \in \mathbb{C} \setminus \mathbb{Z}_0^-). \quad (1.6)$$

where $\mathbb{Z}_0^-$ is the set of non positive integers. The generalized Wright hypergeometric function ${}_p\Psi_q(z)$ is given by the series

$$\begin{aligned} {}_p\Psi_q(z) &= {}_p\Psi_q \left[\begin{matrix} (a_i, \alpha_i)_{1,p} \\ (b_j, \beta_j)_{1,q} \end{matrix} \middle| z \right] \\ &= \frac{\prod_{j=1}^q \Gamma(\beta_j)}{\prod_{i=1}^p \Gamma(\alpha_i)} \sum_{k=0}^{\infty} \frac{\prod_{i=1}^p \Gamma(a_i + \alpha_i k)}{\prod_{j=1}^q \Gamma(b_j + \beta_j k)} \frac{z^k}{k!}, \end{aligned} \quad (1.7)$$

where $a_i, b_j \in \mathbb{C}$, and $\alpha_i, \beta_j \in \mathbb{R}$ ($i = 1, 2, \dots, p; j = 1, 2, \dots, q$). Asymptotic behavior of this function for large values of argument of $z \in \mathbb{C}$ were studied in [17] and under the condition

$$\sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i > -1 \quad (1.8)$$

was found in the work of [18, 19]. Properties of this generalized Wright function were investigated in [21], (see also [22, 23]). In particular, it was proved [21] that ${}_p\Psi_q(z)$, $z \in \mathbb{C}$ is an entire function under the condition (1.8).

If we put $\alpha_1 = \dots = \alpha_p = \beta_1 = \dots = \beta_q = 1$ in (1.7), then the special case of the generalized Wright function is:

$${}_p\Psi_q(z) = {}_p\Psi_q \left[\begin{matrix} (\alpha_1, 1), \dots, (\alpha_p, 1) \\ (\beta_1, 1), \dots, (\beta_q, 1) \end{matrix} \middle| z \right] = \frac{\prod_{j=1}^q \Gamma(\beta_j)}{\prod_{i=1}^p \Gamma(\alpha_i)} {}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p \\ \beta_1, \dots, \beta_q \end{matrix} \middle| z \right] \quad (1.9)$$

provided $0 < \Re(\mu) < \Re(\lambda)$. Here our main aim to establish two generalized integral formulas, which are expressed in terms of generalized Wright functions by inserting generalized Bessel-Maitland function.

The unified integral formulas involving special functions attract the attention many authors in eighteenth century (see, [8],[9]). In 1888, Pincherle studied the integrals involving product of gamma functions along vertical lines (see [2, 3, 4]). The extension of such study carried by Barnes [5], Mellin [6] and Cahen [7] and applied some of these integrals in the study of Riemann zeta function and other Dirichlet's series. The unified integral representation of the Fox H -functions given by Garg [10] and the unified integral representation of the hypergeometric ${}_2F_1$ functions by Ali [11]. Recently, Choi and Agarwal [12] obtained two unified integral representations of Bessel functions $J_\nu(z)$. For more details are found

in various recent works (see [13],[26],[27],[28], [29], [30]). For the present investigation, we need the following result of Oberhettinger [14]

$$\int_0^\infty x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-\lambda} dx = 2\lambda a^{-\lambda} \left(\frac{a}{2} \right)^\mu \frac{\Gamma(2\mu) \Gamma(\lambda - \mu)}{\Gamma(1 + \lambda + \mu)} \quad (1.10)$$

2. MAIN RESULTS

Theorem 1. For $\delta, \lambda, v, c \in \mathbb{C}$, then the following integral formula hold true:

$$\begin{aligned} & \int_0^\infty x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-\lambda} J_{v,q}^{\delta,\gamma} \left(\frac{y}{x + a + \sqrt{x^2 + 2ax}} \right) dx \\ &= \frac{\Gamma(2\mu)}{\Gamma(\gamma)} a^{\mu-\lambda} 2^{1-\mu} \\ & \quad \times {}_3\Psi_3 \left[\begin{matrix} (\gamma, q), (\lambda + 1, 1), (\lambda - \mu, 1); \\ (\nu + 1, \delta), (1 + \lambda + v + \mu, 1), (\lambda, 1); \end{matrix} \quad -\frac{y}{a} \right] \end{aligned}$$

Proof. By applying (1.2) to the LHS of theorem 1 and interchanging the order of integration and summation, which is verified by uniform convergence of the involved series under the given conditions, we obtain

$$\begin{aligned} & \int_0^\infty x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-\lambda} J_{v,q}^{\delta,\gamma} \left(\frac{y}{x + a + \sqrt{x^2 + 2ax}} \right) dx \\ &= \int_0^\infty x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-\lambda} \\ & \quad \times \sum_{n=0}^\infty \frac{(\gamma)_{qn} (-y)^n}{n! \Gamma(\delta n + v + 1)} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-n} dx \\ &= \sum_{n=0}^\infty \frac{(\gamma)_{qn} (-y)^n}{n! \Gamma(\delta n + v + 1)} \int_0^\infty x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-(\lambda+n)} dx \end{aligned}$$

Since $\Re(\lambda) > \Re(\mu) > 0$, $k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

Applying (1.10) to the integrand of theorem 1 and we obtain the following expression:

$$\begin{aligned} &= \sum_{n=0}^\infty \frac{(\gamma)_{qn} (-y)^n}{n! \Gamma(\delta n + v + 1)} 2^{1-\mu} a^{\mu-\lambda} a^{-(\lambda+n)} \left(\frac{a}{2} \right)^\mu \\ & \quad \times \frac{\Gamma(2\mu) \Gamma(\lambda + n - \mu)}{\Gamma(1 + \lambda + \mu + n)} \end{aligned}$$

By making the use of the equation (1.3), we have

$$= 2^{1-\mu} a^{\mu-\lambda} \frac{\Gamma(2\mu)}{\Gamma(\gamma)}$$

$$\begin{aligned} & \times \sum_{k=0}^{\infty} \frac{\Gamma(\gamma + qn) \Gamma(\lambda + n - \mu) \Gamma(\lambda + 1 + n)}{n! \Gamma(\delta n + v + 1) \Gamma(\lambda + n) (1 + \lambda + \mu + n)} \\ & \times \left(-\frac{y}{a}\right)^n \end{aligned}$$

In view of (1.9), we obtain the desired result. $\square$

Corollary 2.1. *If we set $q = 0$ and $\gamma = 1$, then we obtain integral representation of ordinary Bessel-Maitland function (1.1) as:*

$$\begin{aligned} & \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax}\right)^{-\lambda} J_v^{\delta} \left(\frac{y}{x + a + \sqrt{x^2 + 2ax}}\right) dx \\ & = \Gamma(2\mu) a^{\mu-\lambda} 2^{1-\mu} \\ & \quad \times {}_2\Psi_3 \left[\begin{matrix} (\lambda + 1, 1), (\lambda - \mu, 1); \\ (\nu + 1, \delta), (1 + \lambda + v + \mu, 1), (\lambda, 1); \end{matrix} \quad -\frac{y}{a} \right] \end{aligned}$$

Corollary 2.2. *If we set $q = 0$ and $\delta = \gamma = 1$, then we obtain integral representation of ordinary Bessel-Maitland function (1.1) as:*

$$\begin{aligned} & \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax}\right)^{-\lambda} J_v \left(\frac{y}{x + a + \sqrt{x^2 + 2ax}}\right) dx \\ & = \Gamma(2\mu) a^{\mu-\lambda} 2^{1-\mu} \\ & \quad \times {}_2\Psi_3 \left[\begin{matrix} (\lambda + 1, 1), (\lambda - \mu, 1); \\ (\nu + 1, 1), (1 + \lambda + v + \mu, 1), (\lambda, 1); \end{matrix} \quad -\frac{y}{a} \right] \end{aligned}$$

Theorem 2. *For $\delta, \lambda, v, c \in \mathbb{C}$, then the following integral formula hold true:*

$$\begin{aligned} & \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax}\right)^{-\lambda} J_{v,q}^{\delta,\gamma} \left(\frac{xy}{x + a + \sqrt{x^2 + 2ax}}\right) dx \\ & = 2^{1-\mu} a^{\mu-\lambda} \frac{\Gamma(\lambda - \mu)}{\Gamma(\gamma)} \\ & \quad \times {}_3\Psi_3 \left[\begin{matrix} (\gamma, q), (\lambda + 1, 1), (2\mu, 2); \\ (\nu + 1, \delta), (1 + \lambda + v + \mu, 2), (\lambda, 1); \end{matrix} \quad -\frac{y}{a} \right] \end{aligned}$$

Proof. By applying (1.2) to the LHS of theorem 2 and interchanging the order of integration and summation, which is verified by uniform convergence of the involved series under the given conditions, we obtain

$$\begin{aligned} & \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax}\right)^{-\lambda} J_{v,q}^{\delta,\gamma} \left(\frac{xy}{x + a + \sqrt{x^2 + 2ax}}\right) dx \\ & = \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax}\right)^{-\lambda} \end{aligned}$$

$$\begin{aligned}
& \times \sum_{n=0}^{\infty} \frac{(\gamma)_{qn} (-xy)^n}{n! \Gamma(\delta n + v + 1)} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-n} dx \\
& = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn} (-y)^n}{n! \Gamma(\delta n + v + 1)} \int_0^{\infty} x^{\mu+n-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-(\lambda+n)} dx
\end{aligned}$$

Since $\Re(\lambda) > \Re(\mu) > 0$, $k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

Applying (1.10) to the integrand of theorem 2 and we obtain the following expression:

$$\begin{aligned}
& = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn} (-y)^n}{n! \Gamma(\delta n + v + 1)} 2^{\lambda+n} a^{-(\lambda+n)} \left(\frac{a}{2} \right)^{\mu+n} \\
& \quad \times \frac{\Gamma(2(\mu+n)) \Gamma(\lambda+n-\mu-n)}{\Gamma(1+\lambda+\mu+n+n)}
\end{aligned}$$

By making the use of the equation (1.3), we have

$$\begin{aligned}
& = 2^{1-\mu} a^{\mu-\lambda} \frac{\Gamma(\lambda-\mu)}{\Gamma(\gamma)} \\
& \quad \times \sum_{k=0}^{\infty} \frac{\Gamma(\gamma+qn) \Gamma(\lambda+1+n) \Gamma(2\mu+2n)}{n! \Gamma(\delta n + v + 1) \Gamma(\lambda+n) \Gamma(1+\lambda+\mu+2n)} \\
& \quad \times \left(-\frac{y}{2} \right)^n
\end{aligned}$$

In view of (1.9), we obtain the desired result. $\square$

Corollary 2.3. *If we set $q = 0$ and $\gamma = 1$, then we obtain integral representation of ordinary Bessel-Maitland function (1.1) as:*

$$\begin{aligned}
& \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-\lambda} J_v^{\delta} \left(\frac{xy}{x + a + \sqrt{x^2 + 2ax}} \right) dx \\
& = 2^{1-\mu} a^{\mu-\lambda} \Gamma(\lambda-\mu) \\
& \quad \times {}_2\Psi_3 \left[\begin{matrix} (\lambda+1, 1), (2\mu, 2); \\ (\nu+1, \delta), (1+\lambda+v+\mu, 2), (\lambda, 1); \end{matrix} \quad -\frac{y}{a} \right]
\end{aligned}$$

Corollary 2.4. *If we set $q = 0$ and $\delta = \gamma = 1$, then we obtain integral representation of Bessel function $J_v(z)$ as:*

$$\begin{aligned}
& \int_0^{\infty} x^{\mu-1} \left(x + a + \sqrt{x^2 + 2ax} \right)^{-\lambda} J_v \left(\frac{xy}{x + a + \sqrt{x^2 + 2ax}} \right) dx \\
& = 2^{1-\mu} a^{\mu-\lambda} \Gamma(\lambda-\mu) \\
& \quad \times {}_2\Psi_3 \left[\begin{matrix} (\lambda+1, 1), (2\mu, 2); \\ (\nu+1, 1), (1+\lambda+v+\mu, 2), (\lambda, 1); \end{matrix} \quad -\frac{y}{a} \right]
\end{aligned}$$

Conclusion:

Certain unified integral representation of Bessel-Maitland function and its special cases are derived in this study. In this sequel, one can obtain integral representation of more generalized special functions, which has much application in physics and engineering Science.

REFERENCES

- [1] O.I. Marichev, Handbook of Integral Transform and higher transdental functions, Theory and Algorithm tables, Ellis Horwood, Chichester, New York, 1983
- [2] S. Pincherle, Sulle funzioni ipergeometriche generalizzate Nota I, Atti della Reale Accademia dei Lincei. Rendiconti della Classe di Scienze Fisiche, Matematiche e Naturali, Serie 4, Vol. 4 (1888), 694–700
- [3] Pincherle, S: Sulle funzioni ipergeometriche generalizzate. Nota I-II. In: Opere Scelte-I, Unione Matematica Italiana, Ed. Cremonese, Rome (1954), 223-239.
- [4] Mainardi, F, Pagnini, G, Pincherle, S: The pioneer of the Mellin-Barnes integrals. J. Comput. Appl. Math. 153, (2003), 331-342.
- [5] Barnes, EW: A new development of the theory of the hypergeometric functions. Proc. Lond. Math. Soc. 6, 141-177, (1908)
- [6] Mellin, HM: Abriß einer einheitlichen theorie der Gamma und der hypergeometrischen funktionen. Math. Ann. 68, 305-337 (1910)
- [7] Cahen, E: Sur la fonction $\xi\sigma$ de Riemann et sur des fonctions analogues. Ann. Sci. Éc. Norm. Super. 11, 75-164, (1894)
- [8] Brychkov, YA: Handbook of special functions: derivatives, integrals, series and other formulas. CRC Press, Boca Raton, (2008)
- [9] Choi, J, Hasanov, A, Srivastava, HM, Turaev, M: Integral representations for Srivastava's triple hypergeometric functions. Taiwan. J. Math. 15, 2751-2762 (2011)
- [10] Garg, M, Mittal, S: On a new unified integral. Proc. Indian Acad. Sci. Math. Sci. 114(2), 99-101 (2003)
- [11] Ali, S: On some new unified integrals. Adv. Comput. Math. Appl. 1(3), 151-153 (2012)
- [12] Junesang Choi, Praveen Agarwal, Certain unified integrals associated with Bessel functions, Boundary Value Problems 2013, 2013:95
- [13] J.Choi, P.Agarwal, S.Mathur and S.D.Purohit, Certain new integral formulas involving the generalized Bessel functions, Bull. Korean Math. Soc., 4(51), 2014, 995-1003.
- [14] Oberhettinger, F: Tables of Mellin Transforms. Springer, New York (1974)
- [15] Watson, GN: A Treatise on the Theory of Bessel Functions. Cambridge Mathematical Library Edition. Cambridge University Press, Cambridge (1995). Reprinted (1996)
- [16] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher transcendental functions. Vols. I, II*, McGraw-Hill Book Company, Inc., New York, 1953. New York, Toronto, London, 1953.
- [17] C. Fox, The asymptotic expansion of generalized hypergeometric functions, Proc. London. Math. Soc. **27** (1928), no. 4, 389-400.

- [18] E. M. Wright, The asymptotic expansion of integral functions defined by Taylor series, Philos. Trans. Roy. Soc. London, Ser. A. **238** (1940), 423–451.
- [19] E. M. Wright, The asymptotic expansion of the generalized hypergeometric function, Proc. London Math. Soc. (2) **46** (1940), 389–408.
- [20] A. A. Kilbas and M. Saigo, *H-transforms*, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [21] A. A. Kilbas, M. Saigo and J. J. Trujillo, On the generalized Wright function, Fract. Calc. Appl. Anal. **5** (2002), no. 4, 437–460.
- [22] A. A. Kilbas and N. Sebastian, Generalized fractional integration of Bessel function of the first kind, Integral Transforms Spec. Funct. **19** (2008), no. 11–12, 869–883.
- [23] A. A. Kilbas and N. Sebastian, Fractional integration of the product of Bessel function of the first kind, Fract. Calc. Appl. Anal. **13** (2010), no. 2, 159–175.
- [24] E. D. Rainville, *Special functions*, Macmillan, New York, 1960.
- [25] M. Singh, M.A. Khan and A.H. Khan, On some properties of a generalization of Bessel-Maitland function, International journal of Mathematics trends and technology, 14(1), 2014
- [26] K.S. Nisar, R. K. Parmar, and A.H. Abusufian, Certain New Unified Integrals Associated With The Generalized K-Bessel Function, Far East Journal of Mathematical Sciences 2016.
- [27] K. S. Nisar and S. R. Mondal, Certain unified integral formulas involving the generalized modified k -Bessel function of the first kind, arXiv:1601.06487 [math.CA].
- [28] K. S. Nisar, P. Agarwal and S. Jain, Some unified integrals associated with Bessel-Struve kernel function, arXiv:1602.01496v1 [math.CA]
- [29] N. Menaria , D. Baleanu and S. D. Purohit, Integral Formulas Involving Product of General Class of Polynomials and Generalized Bessel Function, Sohag J. Math. 3, No. 2, 1-5 (2016)
- [30] N. Menaria, S. D. Purohit and R. K. Parmar, On a new class of integrals involving generalized Mittag-leffler function, Surveys in Mathematics and its Applications, Volume 11 (2016), 1-9.

M.S. ABOUZAIID: DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KAFRELSHIEKH UNIVERSITY, EGYPT.

E-mail address: moheb_abouzaid@hotmail.com

A.H. ABUSUFIAN: ¹COLLEGE OF ARTS AND SCIENCE, PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY, WADI AL DAWASER, RIYADH REGION 11991, SAUDI ARABIA

E-mail address: sufianmath97@hotmail.com

K. S. NISAR: DEPARTMENT OF MATHEMATICS, COLLEGE OF ARTS AND SCIENCE, PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY, WADI AL DAWASER, RIYADH REGION 11991, SAUDI ARABIA

E-mail address: ksnisar1@gmail.com